



## F10 gene

coagulation factor X

### Normal Function

The *F10* gene provides instructions for making a protein called coagulation factor X. Coagulation factors are a group of related proteins that are involved in the coagulation system, which is a series of chemical reactions that form blood clots. After an injury, clots seal off blood vessels to stop bleeding and trigger blood vessel repair.

Coagulation factor X is made primarily by cells in the liver. The protein circulates in the bloodstream in an inactive form until the coagulation system is turned on (activated) by an injury that damages blood vessels. When coagulation factor X is activated, it interacts with other coagulation factors to convert an important coagulation protein called prothrombin to its active form, thrombin. Thrombin then converts a protein called fibrinogen into fibrin, which is the material that forms blood clots.

### Health Conditions Related to Genetic Changes


#### factor X deficiency

At least 130 mutations in the *F10* gene have been found to cause a rare bleeding disorder called factor X deficiency. This disorder commonly causes nosebleeds, easy bruising, bleeding under the skin, bleeding of the gums, blood in the urine (hematuria), and prolonged or excessive bleeding following surgery or trauma. Some *F10* gene mutations that cause factor X deficiency reduce the amount of coagulation factor X in the bloodstream, resulting in a form of the disorder called type I. Other *F10* gene mutations result in the production of a coagulation factor X protein with impaired function, leading to type II factor X deficiency. Reduced quantity or function of coagulation factor X prevents blood from clotting normally, causing episodes of abnormal bleeding that can be severe.

## Chromosomal Location

Cytogenetic Location: 13q34, which is the long (q) arm of chromosome 13 at position 34

Molecular Location: base pairs 113,122,799 to 113,149,529 on chromosome 13 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

## Other Names for This Gene

- autoprothrombin III
- prothrombinase
- Prower factor
- Stuart factor
- Stuart-Prower factor

## Additional Information & Resources

### Educational Resources

- Biochemistry (fifth edition, 2002): Blood-Clotting Cascade (image)  
<https://www.ncbi.nlm.nih.gov/books/NBK22589/?rendertype=figure&id=A1401>

### Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28coagulation+factor+X%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1080+days%22%5Bdp%5D>

### OMIM

- COAGULATION FACTOR X  
<http://omim.org/entry/613872>

## Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology  
[http://atlasgeneticsoncology.org/Genes/GC\\_F10.html](http://atlasgeneticsoncology.org/Genes/GC_F10.html)
- ClinVar  
<https://www.ncbi.nlm.nih.gov/clinvar?term=F10%5Bgene%5D>
- HGNC Gene Family: Gla domain containing  
<http://www.genenames.org/cgi-bin/genefamilies/set/1250>
- HGNC Gene Symbol Report  
[http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?q=data/hgnc\\_data.php&hgnc\\_id=3528](http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=3528)
- NCBI Gene  
<https://www.ncbi.nlm.nih.gov/gene/2159>
- UniProt  
<http://www.uniprot.org/uniprot/P00742>

## **Sources for This Summary**

- Brown DL, Kouides PA. Diagnosis and treatment of inherited factor X deficiency. *Haemophilia*. 2008 Nov;14(6):1176-82. doi: 10.1111/j.1365-2516.2008.01856.x.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/19141158>
- OMIM: COAGULATION FACTOR X  
<http://omim.org/entry/613872>
- Girolami A, Scarparo P, Scandellari R, Allemand E. Congenital factor X deficiencies with a defect only or predominantly in the extrinsic or in the intrinsic system: a critical evaluation. *Am J Hematol*. 2008 Aug;83(8):668-71. doi: 10.1002/ajh.21207. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/18506702>
- Girolami A, Vettore S, Scarparo P, Lombardi AM. Persistent validity of a classification of congenital factor X defects based on clotting, chromogenic and immunological assays even in the molecular biology era. *Haemophilia*. 2011 Jan;17(1):17-20. doi: 10.1111/j.1365-2516.2010.02328.x.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/20546029>
- Herrmann FH, Auerswald G, Ruiz-Saez A, Navarrete M, Pollmann H, Lopaciuk S, Batorova A, Wulff K; Greifswald Factor X Deficiency Study Group. Factor X deficiency: clinical manifestation of 102 subjects from Europe and Latin America with mutations in the factor 10 gene. *Haemophilia*. 2006 Sep;12(5):479-89.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/16919077>
- Karimi M, Menegatti M, Afrasiabi A, Sarikhani S, Peyvandi F. Phenotype and genotype report on homozygous and heterozygous patients with congenital factor X deficiency. *Haematologica*. 2008 Jun;93(6):934-8. doi: 10.3324/haematol.12211. Epub 2008 Apr 9.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/18403394>

- Menegatti M, Peyvandi F. Factor X deficiency. *Semin Thromb Hemost*. 2009 Jun;35(4):407-15. doi: 10.1055/s-0029-1225763. Epub 2009 Jul 13. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/19598069>
  - Uprichard J, Perry DJ. Factor X deficiency. *Blood Rev*. 2002 Jun;16(2):97-110. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/12127953>
- 

Reprinted from Genetics Home Reference:  
<https://ghr.nlm.nih.gov/gene/F10>

Reviewed: January 2015  
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications  
U.S. National Library of Medicine  
National Institutes of Health  
Department of Health & Human Services